

NOMINA MUTANDA.

BY ALBERT COLLIN & E. HARTERT.

1. *Alcippe poiocephala blythi* nom. nov. versus *A. p. magnirostris* Walden.

Alcippe magnirostris Walden, *Blyth's Cat. Mamm. & B. Burma*, p. 115 (1875—Karennee Hills), nec Moore, *Proc. Zool. Soc. London*, 1854, p. 277 (Malacca).

(Specimens from Bandon in the Northern, Siamese, part of the Malay Peninsula show only partially indications of the blackish side-stripes on the crown; a larger series might serve to separate them again.)

2. *Astur fasciatus vigilax* versus *A. f. insularis* Sarasin.

Astur fasciatus insularis Sarasin, *Nova Caledonia, Zool., Av.* p. 8 (1913), nec *Astur insularis* Madarász, *Orn. Monatsber.* xviii, p. 65 (1910—Ceylon).

Astur fasciatus vigilax Wetmore, *Condor* xxviii, p. 46 (1926).

3. *Anas leucophrys* Vieill. versus *A. torquata* Vieill.

Anas torquata Vieillot, *Nouv. Dict. d'Hist. Nat.* v, p. 110 (1816—Paraguay), nec S. G. Gmelin, *Reise d. Russl.* ii, p. 179, pl. xiv (1774).

Anas leucophrys Vieillot, *Nouv. Dict. d'Hist. Nat.* v, p. 156 (1816—Paraguay).

4. *Anthus nicholsoni neumannianus* nom. nov. versus *A. n. longirostris* Neum.

Anthus nicholsoni longirostris Neumann, *Orn. Monatsber.* xiii, p. 77 (1905—"Gardulla am Gandjuli See"), nec *Anthus obscurus longirostris* Brehm, *Naumannia*, v i, p. 342 (1856).

5. *Collocalia esculenta erwini* nom. nov. versus *C. e. maxima* Grant.

Bull. B.O. Club, xxv, p. 35 (1914—Utakwa River), nec *C. maxima* Hume, *Stray Feath.* iv, p. 223 (nomen nudum), vi, p. 49 (1878—Mergui. Not nomen nudum but synonym of *innominata*).

6. *Otus hartlaubi* (Gieb.) versus *Otus leucopsis* (Hartl.).

Athene leucopsis Hartlaub, *Rev. Zool.* 1849, p. 496 (St. Thomé), nec Gould, *Proc. Zool. Soc. London*, 1837, p. 99 (van Diemen's Land).

Noctua hartlaubi Giebel, *Thes. Orn.* ii, p. 717, nom. nov. (1875).

7. *Carduelis ictericus bavarici* nom. nov. versus *Carduelis ictericus campestris* (Spix).

Fringilla campestris Spix, *Av. Bras.* ii, p. 48, pl. lxi, fig. 3 (1825), nec Schrank, *Fauna Boica*, p. 181 (1789—Bei Neuburg, Bayern).

8. *Eopsaltria australis griseogularis* Gould versus *E. a. gularis* (Quoy et Gaim.).

Muscicapa gularis Quoy et Gaimard, *Voy. de l'Australie, Zool.* i, p. 176 (1830—King George Sound, W. Austr.), nec Temminck, *Pl. Col.* 167, fig. 1 (1823—Brazil).

Eopsaltria griseogularis Gould, *Synops. B. Austr.* iv, app., p. 2 (1838—Swan River, W. Australia).

9. *Stoparola thalassina thalassina* (Swains.) versus *S. melanops melanops* (Vig.).

Muscicapa melanops Vigors, *Proc. Zool. Soc. London*, 1831, p. 171 (Himalaya), nec Vieillot, *Nouv. Dict. d'Hist. Nat.* xxi, p. 452 (1818—Paraguay).

Muscicapa thalassina Swainson, *Natur. Libr.* x, Flycatchers, p. 252 (1838—India).

10. **Paroaria rubrifacies** nom. nov. versus *P. cucullata* (Lath.).

Loxia cucullata Latham, *Ind. Orn.* i, p. 378 (1790—Brazil), nec Boddaert, *Tabl. Pl. Enl.* p. 24 (1783—Ex Buffon).

11. **Spermospiza haematina pustulata** (Voigt) versus *S. h. guttata* (Vieill.).

Loxia guttata Vieillot, *Ois. Chant.* p. 103, pl. 68 (1856—Congo), nec Shaw, *Mus. Lever.* ii, p. 47 (1796—"Australia").

Fringilla pustulata Voigt, *Cuvier's Thierreich*, i, p. 581 (1831—Congo).

12. **Monasa atra** (Bodd.) versus *M. nigra* (P. L. S. Müll.).

Cuculus niger P. L. S. Müller, *Natursyst. Suppl.* p. 90 (1776—Cayenne), nec Linnaeus, *Syst. Nat.* ed. x, p. 111 (1758—Bengal).

Cuculus ater Boddaert, *Table Pl. Enl.* p. 30 (1783—Cayenne).

13. **Cyanoderma labuanensis** nom. nov. versus *C. bicolor* (Blyth).

Timalia bicolor Blyth, *Ibis* 1865, p. 46 ("Prepared like the Malaccan specimens," locality therefore unknown, but certainly not from Malacca), nec Lafresnaye, *Mag. de Zool.* 1835, pl. xxxix text.

This form is not rare in Labuan, North Borneo, and restricted to the island of Borneo.

14. **Buteo buteo burmanicus** Oates versus *B. b. japonicus* (Temm. & Schleg.).

Falco buteo japonicus Temminck & Schlegel, Siebold's *Fauna Japon., Aves*, p. 16, Tab. vi (1844—Japan), nec *Falco tinunculus japonicus*, t.c. p. 2, Tab. i (1844).

Buteo burmanicus Oates, *Stray Feath.* iii, p. 30 (1875—Upper Pegu).

15. **Crimiger balicus bartelsi** nom. nov. versus *C. gularis gularis* (Horsf.).

Turdus gularis Horsfield, *Trans. Linn. Soc. London*, xiii, p. 150 (1822—Java), nec Latham, *Ind. Orn. Suppl.* p. xl (1801—Cumberland).

16. **Centropus superciliosus intermedius** Someren.

Bull. B.O. Club, xli, p. 125 (1921—Mombasa) is preoccupied by *Centrocoeryx intermedius* Hume 1873 = *Centropus sinensis intermedius*.

If confirmed as different, a new name must be invented for van Someren's *intermedius*; cf. *NOV. ZOOLOG.* 1925, p. 153.

17. **Monticola semicastanea** nom. nov. versus *M. erythrogaster* (Vig.).

Turdus erythrogaster Vigors, *Proc. Zool. Soc. London*, 1831, p. 171 (Himalaya), nec Boddaert, *Table Pl. Enl.* p. 22 (1783), quid est *Spreo pulcher* (P. L. S. Müll.).

18. **Tyto alba tuidara** (Griff. & Pidgeon) versus *T. a. perlata* (Licht.).

Strix perlata Lichtenstein, *Verz. Doubl.* p. 59 (1823), nec Vieillot, *Nouv. Dict. d'Hist. Nat.* vii, p. 26 (1817).

Strix tuidara Griffith & Pidgeon, *Cuvier's Anim. Kingd., Aves* i, p. 74 (1829—Brazil).

19. **Uroleuca cristatella** (Temm.) versus *U. cyanoleuca* (Wied.).

Corvus cyanoleucus Wied, *Reise Brasil.* ii, p. 190 (1821—Brazil), nec Latham, *Ind. Orn. Suppl.* p. xxv (1801—N.S. Wales).

Corvus cristatellus Temminck, *Pl. Col.* 193 (1822—Brazil).

20. **Gerygone neglecta dohertyi** R. & H. versus *G. n. virescens* (Blyth).

Sylvia virescens (Müller MS.) Blyth, *Ibis*, 1870, p. 169 (New Guinea), nec Vieillot, *Ois. Am. Sept.* ii, p. 42 (1807).

Gerygone neglecta dohertyi Rothschild & Hartert, *Nov. Zool.* 1903, p. 473 (Kapaur, New Guinea).

21. *Mimus plumbeus* nom. nov. versus *Mimus lividus* (Licht.).

Turdus lividus Lichtenstein, *Verz. Doubl.* p. 39 (1823), nec Wilson, *Am. Orn.* ii, p. 90, pl. xiv, fig. 3 (1810).

Turdus cinereus Voigt, Cuvier's *Thierreich*, i, p. 483 (1831), nec Gmelin, *Syst. Nat.* i, 2, p. 810 (1789).

22. *Bradornis pallida bowdleri* nom. nov. versus *B. p. sharpei* R.

Bradyornis pallidus sharpei Rothschild, *Bull. B.O. Club*, xxxiii, p. 66 (1913—Abyssinia), nec *Bradyornis sharpei* Bocage, *Bull. B.O. Club*, iii, p. xliii (1894—Galanga in Angola).

23. *Anthus spinoletta reuteri* Munst. versus *A. s. borealis* Hesse.

Anthus borealis Hesse, *Journ. f. Orn.* 1915, p. 386 (Sachalin), nec *A. pratensis* var. *borealis* (ex E. v. Homeyer MS.) Blasius, "Naumann, *Naturg. Vög. Mitteleuropas*" (sic!) iii, p. 58 (1900—Anklam in Pommern).

Anthus spinoletta reuteri Munsterhjelm, *Nyt. Mag. for Naturvidensk.* 1916, p. 165 (Sachalin).

24. *Anthus rufogularis* Brehm versus *A. cervinus* Pall.

Anthus rufogularis Brehm, *Lehrb. Nat. eur. Vög.* ii, p. 963 (1824—Nubien und Deutschland).

Motacilla cervina Pallas, *Zoogr. Rosso-Asiat.* i, p. 511 (1827).

25. ? *Fringilla coelebs bellicosa* Floer. versus *F. c. tristis* Floer. ?

Fringilla coelebs tristis Floericke, *Mitt. Österr. Reichsb. f. Vogelk. u. Vogelschutz*, iii, p. 21 (1901—"im Winter auf der kurischen Nehrung"), nec *Fringilla tristis* Linnaeus, *Syst. Nat.* ed. x, p. 181. (1758—"Habitat in America septentrionali").

Fringilla coelebs bellicosus Floericke, *Mitt. Vogelwelt*, xix, p. 105 (1921—Wolhynien).

Fringilla coelebs karelica Räsänen, *Luonnon Ystävä (Naturfreund)*, xxviii, p. 21 (1924—Karelische Halbinsel).

If a north-eastern subspecies of *F. coelebs* can be distinguished, all three above names may perhaps refer to the same form. The descriptions of "*bellicosus*" and "*tristis*," however, do not agree!

26. *Falco peregrinus perconfusus* nom. nov. versus *Falco peregrinus minor* Schleg.

Falco minor Schlegel, *Abh. Geb. Zool. & vergl. Anat.* (2), iii, p. 20 (1844—Kap der Guten Hoffnung), nec *Falco nisus minor* Bekker, Borkhausen & Lichthammer, in Bekker & Lembke, *Teutsche Orn.*, iii, pl. 1-5 (1800-1811, Deutschland).

Falco peregrinus var. *capensis* Grill, *Sr. vet. ak. handl.* ii, 10, p. 48 (1858), nec *F. capensis* Shaw, *Gen. Zool.* vii, p. 192 (1809).

27. *Prunella fulvescens sushkini* nom. nov. versus *P. fulvescens tibetana* Sushk.

Prunella fulvescens tibetana Sushkin, *Proc. Bost. Soc. Nat. Hist.* xxxviii, i, p. 53 (1925), nec *Accentor collaris tibetanus* Bianchi, *Ann. Mus. Zool. Petersburg*, ix, p. 128 (1924) = *Prunella collaris tibetana* (Bianchi).

28. *Lanius collurioides delacouri* nom. nov. versus *L. c. melanocephalus* Delacour.

Bull. B.O. Club, xlvii, p. 13 (1926—Annam), nec *Lanius melanocephalus* Gmelin, *Syst. Nat.* i, 1, p. 309 (1788).

29. *Turdus torquatus caucasicus* nom. nov. versus *T. torquatus orientalis* Seebohm.

Ibis, 1888, p. 311 ("Caucasus and Persia"), nec *Turdus orientalis* Gmelin, *Syst. Nat.* i, 2, p. 821 (1789—India).

30. *Turdus gouldi cinereiceps* nom. nov. versus *T. castaneus castaneus* (Gould).

Merula castanea Gould, *Proc. Zool. Soc. London*, 1835, p. 185 (Himalaya), nec *Turdus castaneus* P. L. S. Müller, *Natursyst. Suppl.* p. 143 (1776).

31. *Alseonax cinereus nigrorum* nom. nov. versus *A. cinereus cinerascens* (Sharpe).

Muscicapa cinerascens Sharpe, *Cat. B. Brit. Mus.* iv, p. 155 (1879—Gold Coast), nec Spix, *Aves Bras.* ii, p. 16, pl. xii (1825). Cf. Bates, *Ibis*, 1926, p. 584.

UG 1927
BASED

